

INSUFFICIENCY DYSBIOSIS

Insufficiency dysbiosis is characterized by low levels of beneficial bacteria that provide critical support for healthy intestinal and immune function. Insufficient levels of beneficial bacteria may result in an elevated risk of intestinal infections, increased intestinal permeability, decreased protective factors such as secretory IgA, and increased inflammation.

MARKERS CHARACTERIZING INSUFFICIENCY DYSBIOSIS

Normal Bacterial Flora: <i>(low levels)</i>	<i>Bacteroides fragilis</i> <i>Bifidobacterium</i> spp. <i>Enterococcus</i> spp. <i>Escherichia</i> spp. <i>Lactobacillus</i> spp. <i>Akkermansia mucinophilia</i> <i>Faecalibacterium prausnitzii</i>
Phyla Microbiota: <i>(low levels)</i>	<i>Bacteroidetes</i> <i>Firmicutes</i>
Associated Intestinal Health Markers:	Secretory IgA <i>(often low to very low levels)</i> Anti-gliadin IgA <i>(often elevated)</i> Zonulin <i>(sometimes elevated)</i>

Continued...

INFLAMMATORY DYSBIOSIS

Inflammatory dysbiosis is characterized by moderate to high levels of certain pathogens, normal microbiota, and opportunistic microbes that promote inflammation and increased intestinal permeability. Many pro-inflammatory microbes are gram-negative bacteria that belong to the Proteobacteria phylum and produce a form of lipopolysaccharide (*LPS*) that is a potent activator of inflammatory responses.

MARKERS CHARACTERIZING INFLAMMATORY DYSBIOSIS	
Pathogens (low to high levels)	<i>Campylobacter</i> <i>C. difficile</i> Pathogenic <i>E. coli</i> <i>Salmonella</i> <i>Vibrio cholerae</i> <i>Yersinia enterocolitica</i> <i>Giardia</i>
Normal Flora (high levels)	<i>Escherichia</i> spp. <i>Enterobacter</i> spp.
Opportunistic Bacteria, Yeast and Protozoa (moderate to high levels)	<i>Morganella</i> spp. <i>Pseudomonas</i> spp. <i>Pseudomonas aeruginosa</i> <i>Citrobacter</i> spp. <i>Citrobacter freundii</i> <i>Klebsiella</i> spp. <i>Klebsiella pneumoniae</i> <i>Proteus</i> spp. <i>Proteus mirabilis</i> <i>Fusobacterium</i> spp. <i>Candida</i> spp. <i>Candida albicans</i> Parasitic protozoa
Associated Intestinal Health Markers:	Secretory IgA (often low levels, but sometimes elevated) Calprotectin (often elevated, but sometimes very low levels) Zonulin (may be elevated in some cases)

Continued...



GI Microbial Assay Plus

DIGESTIVE DYSFUNCTION DYSBIOSIS

Dysbiosis associated with digestive dysfunction is very common, and is often due to low stomach acid (hypochlorhydria), insufficient bile acids, poor digestion (*pancreatic insufficiency or brush border enzyme deficiency*), reduced absorption, and altered gastrointestinal motility. Altered digestion and motility can result in imbalances in the microbiome, characterized by overgrowth of certain species.

MARKERS ASSOCIATED WITH DIGESTIVE DYSFUNCTION	
Pathogens (low to high levels)	Most types, especially if multiple pathogens are present
<i>H. pylori</i> (moderate to high levels)	<i>Helicobacter pylori</i> (with or without virulence factors)
Normal Flora (high levels)	<i>Enterococcus</i> <i>Lactobacillus</i> <i>Clostridium</i>
Phyla Microbiota (high levels)	<i>Bacteroidetes</i> and/or <i>Firmicutes</i>
Opportunistic Bacteria, Yeast and Parasites (moderate to high levels)	<i>Bacillus</i> spp. <i>Enterococcus faecalis</i> <i>Enterococcus faecium</i> <i>Morganella</i> spp. <i>Staphylococcus</i> spp. <i>Staphylococcus aureus</i> <i>Streptococcus</i> spp. <i>Methanobacteriaceae</i> (family) <i>Klebsiella pneumoniae</i> <i>Prevotella copri</i> <i>Candida</i> spp. <i>Candida albicans</i> <i>Parasitic Protozoa</i>
Associated Intestinal Health Markers:	Elastase-1 (often low to moderately low levels) Steatocrit (sometimes elevated)
Associated Symptoms & Conditions:	Excessive gas and bloating Stomach or abdominal discomfort Feeling of fullness after a meal Heartburn, gastroesophageal reflux Constipation or diarrhea Small intestinal bacterial overgrowth (SIBO) Food sensitivities and intolerances



GI Microbial Assay Plus