

5895 Shiloh Rd, Ste 101 | Alpharetta GA 30005 877.485.5336

Patient: Ima Sample

Collected: 6/1/2025 DOB: 2/1/1980 Accession: 00000000-0000

Received: 6/2/2025 Completed: 6/15/2025

Ordered by: Sample Provider, MD

DNA STOOL ANALYSIS BY QUANTITATIVE PCR

YOUR PERSONALIZED REPORT

PATHOGENS

The GI-MAP® includes pathogens (bacterial, parasitic and viral) commonly known to cause gastroenteritis. Note that not all individuals with positive findings will present with symptoms. Many factors, including the health of the individual (such as immune health, digestive function, and microbiome balance), the transient nature of most pathogens, and the presence and expression of virulence factors, all contribute to pathogen virulence and individual symptoms.

BACTERIAL PATHOGENS	Result	Reference
Campylobacter	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
C. difficile Toxin A	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
C. difficile Toxin B	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
E. coli - EPEC/EHEC	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
E. coli O157	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
Enteroinvasive E. coli/Shigella	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
Enterotoxigenic E. coli LT/ST	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
Shiga-like Toxin <i>E. coli</i> stx1	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
Shiga-like Toxin <i>E. coli</i> stx2	<dl< td=""><td>< 1.00e3</td></dl<>	< 1.00e3
Salmonella	2.04e3	< 1.00e4
Vibrio cholerae	<dl< td=""><td>< 1.00e5</td></dl<>	< 1.00e5
Yersinia enterocolitica	<dl< td=""><td>< 1.00e5</td></dl<>	< 1.00e5
PARASITIC PATHOGENS		
Cryptosporidium	<dl< td=""><td>< 1.00e6</td></dl<>	< 1.00e6
Entamoeba histolytica	<dl< td=""><td>< 1.00e4</td></dl<>	< 1.00e4
Giardia	<dl< td=""><td>< 5.00e3</td></dl<>	< 5.00e3
VIRAL PATHOGENS		
Adenovirus 40/41	<dl< td=""><td>< 1.00e10</td></dl<>	< 1.00e10
Norovirus GI/II	<dl< td=""><td>< 1.00e7</td></dl<>	< 1.00e7

KEY: Results are reported as genome equivalents per gram of stool, which is a standard method for estimating the number of microbes measured per gram of stool, based on qPCR analysis of DNA samples.

Results are expressed in standard scientific notation. For example, a reported result of 3.5e7 is equivalent to 3.5 x 10⁷ microbes per gram, which equals 35,000,000 (35 million) microbes per gram of stool.

< dl represents results below detectable limit.

Patient: Sample, Ima Accession: 00000000-0000

HELICOBACTER PYLORI			
H. PYLORI & VIRULENCE FACTORS	Result	Reference	
Helicobacter pylori	8.18e2	< 1.00e3	
Virulence Factor, babA	Negative	Negative	
Virulence Factor, cagA	Negative	Negative	
Virulence Factor, dupA	Negative	Negative	
Virulence Factor, iceA	Negative	Negative	
Virulence Factor, oipA	Negative	Negative	
Virulence Factor, vacA	Negative	Negative	
Virulence Factor, virB	Negative	Negative	
Virulence Factor, virD	Negative	Negative	
	Nogalivo	14094440	

Result Reference Amoxicillin Negative Negative Genes associated with amoxicillin resistance PBP1A S414R Absent PBP1A T556S Absent PBP1A N562Y Absent Result Reference Result Reference Result Reference Reference Reference Clarithromycin Genes associated with clarithromycin A2142C A2142G A2143G

	Result	Reference
Fluoroquinolones	Positive	Negative
Genes associated with fluoroquinolone resistance		
gyrA N87K	Present	
gyrA D91G	Present	
gyrA D91N	Present	
gyrB S479N	Absent	
gyrB R484K	Absent	

	Result	Reference
Clarithromycin	Positive	Negative
Genes associated with clarithromycin resistance		
A2142C	Present	
A2142G	Present	
A2143G	Present	

	Result	Reference
Tetracycline	Negative	Negative
Genes associated with tetracycline resistance		
A926G	Absent	
AGA926-928TTC	Absent	